

- Buy in bulk and bring your own cloth or recycled grocery bags to the store
- Keep litter, leaves and debris out of the street gutters and storm drains
- Re-use when possible
- Reduce consumption by avoiding excessively packaged products
- Use environmentally friendly cleaners or products that are low in phosphorous to reduce the amount of nutrients discharged into our lakes, streams and coastal waters
- Choose products packaged in recycled materials

Join the Algalita Marine Research Foundation (AMRF)

AMRF is a 501c(3) nonprofit organization founded in Long Beach, California and is dedicated to the protection of the marine environment through research and education. Our primary work is to establish a baseline data set of the level of plastic debris found in our oceans and inform the public about its existence through education and public education.

Membership Levels

- Basic membership** \$25
- Friend of Algalita** \$50
- Good Friend of Algalita** \$100
- Great Friend of Algalita** \$150

A membership donation of \$500 or more will receive an AMRF t-shirt and a "Research Clips" video.

Member Information:

Name _____

Address _____

Phone # _____

e-mail _____

Send by mail or fax.

Algalita Marine Research Foundation

148 North Marina Drive, Long Beach, California 90803
 Phone: (562) 598-4889 Fax: (562) 598-0712
 e-mail: info@algalita.org or sign up online.

www.algalita.org
 www.plasticdebris.org



We drink out of them, eat off of them, sit on them and even drive in them. They are durable, lightweight, and can be made into virtually anything. But it is these useful properties of plastics which make them so harmful when they end up in the environment.

Plastics, like diamonds... ARE FOREVER!

- Only 3.5% of plastics are recycled in any way
- 63 pounds of plastic packaging goes into landfills in the U.S. per person per year.
- Broken, degraded plastic pieces outweigh surface zooplankton in the Central North Pacific by 6 to 1.

Why is plastic in the ocean a problem?

Because plastic does not biodegrade. When something biodegrades, naturally occurring organisms break down natural materials into their simple chemical components. Paper, when it breaks down, becomes carbon dioxide and water. But plastic, a synthetic material, never biodegrades. Instead, plastic goes through a process called photodegradation, where it is broken down by sunlight into smaller and smaller pieces, all of which are still plastic polymers. Even this degradation process can take a very long time. Estimates of 500 years for a disposable diaper, 400 years for a plastic six-pack ring and 450 years for a plastic bottle have been made. The more plastic we produce, the more we have to live with...forever!

The ocean is especially susceptible to plastic pollution, because:

1. It takes longer for the sun to break apart a piece of plastic in the ocean than a piece of plastic on land. This is because the ocean water cools the plastic piece, prevents heat build-up and limits UV light exposure.
2. Plastics are carried by currents, which can concentrate the plastic in certain areas and prevent it from washing onshore. Circulating currents in the ocean caused by stable weather patterns are called "gyres." When plastic is flushed out of a gyre by storms and washes ashore, or when rain sends plastics down rivers to the sea, much of it is mixed into beach sand and can never be recovered.



Some fish mistake plastic "nurdles" for food

But plastic pollution doesn't just look bad...

Plastic pollution is bad for the millions of animals that inhabit our ocean waters and for the people who fish, swim and recreate there.

Many marine birds and animals mistakenly eat pelagic (free-floating) plastic.

- Often these animals cannot distinguish plastic from food. Plastic, because of its high molecular weight and the nature of its chemical bonds, can never be digested. It provides no

nutrients. Eating plastic can cause animals to feel full and not hungry even though they are not actually consuming food. In birds, it has been shown that ingestion of plastics can prevent migration and reproduction, and can eventually cause starvation and death. In turtles, plastic has been shown to block intestines and make the animals float so that they cannot dive for food.

- Toxic chemicals in plastics can make marine birds and animals sick. Over 80 species of seabirds have been found to ingest plastic. Sea bird chicks are especially vulnerable as they receive high levels of pollution from the yolk sac and, after hatching, from food brought by their parents.



A plastic bottle takes 450 years to degrade.

- Ninety percent of Laysan Albatross chick carcasses and regurgitated food boluses contain plastic.
- Marine birds and animals can become entangled in plastic nets and fishing line. An estimated 100,000 marine mammal deaths occur this way each year in the North Pacific.
- Chemicals used to make plastics can escape into the atmosphere during the manufacturing process. Fourteen percent of the toxic airborne chemicals nationally are from "plastics sector" releases. These chemicals can be toxic or carcinogenic, harming both people and animals.

The plastic future looks grim

Virgin plastic pellets are released into the environment by thousands of consumer plastics manufacturers and are the most common contaminant on some beaches.

Scientists predict a 10-fold increase in ocean plastics by the year 2010, which would bring the ratio of surface plastic to zooplankton in the North Pacific Central Gyre to 60:1 by weight.

What are the facts about plastic?

The magnitude of our plastic problem is enormous.

- The American people weigh approximately 50 billion pounds, but 100 billion pounds of plastic resin pellets (the raw materials for consumer plastics) are produced in the U.S. annually.
- 63 pounds of plastic packaging goes to landfills in the U.S. per person per year.



Only 3.5% of plastics are recycled in any way.

- Only 3.5% of plastic is recycled in any way. Reheating plastic gives it a “heat history” which reduces its flexibility. Reheating temperatures are too low to burn off contaminants; therefore, very few plastics are recycled into the same type of container or product that they were originally. Usually, recycled merely means collected, not reprocessed into useful products.
- The “chasing arrows” symbol (shown above) only denotes type of plastic; otherwise it is meaningless. Plastics manufacturers adopted the symbol over the protests of environmentalists and are now being challenged in court by several cities over its implications.

How are plastics getting in the ocean?

Approximately 100 million containers are shipped annually over the world’s oceans. Shipping across the North Pacific Ocean from Asia to North America is along Great Circle routes in the West Wind Drift current at the northern edge of the Central Pacific Gyre. Frequent severe storms along this route cause the loss of hundreds of containers overboard each year contributing, among other plastics, tens of thousands of shoes and millions of plastic shopping bags made in Asia.

Only about 20% of ocean pollution comes from activities at sea. Activities on land contribute most of the remaining 80%.

Because of their buoyancy and persistence, plastic items contribute disproportionately to the overall impact of marine debris. Most of the debris that either entangles animals or is found in their stomachs is made of plastic.

The majority of the plastic that ends up in the Central Pacific Gyre (an area the size of Texas) has been shown to circulate there for at least twelve years. Debris lost in the Bering Sea or the western portion of the Subarctic Gyre will end up there in 3 to 6 years.

Not all plastic floats on the surface. Approximately half of plastics are negatively buoyant. They therefore do not receive sunlight to facilitate the photodegradation process that breaks them into smaller pieces. This debris accumulates on the bottom of the ocean or “benthos.” There the particulates are used by polychaete worms to make their dwelling tubes and some are eaten by flounders and lobsters. The nets, traps and lines that do not photodegrade continue to “ghost fish” (catch fish without a fisherman) and entangle fish and mammals for years.

SIX MISCONCEPTIONS ABOUT PLASTIC

promoted by two powerful plastic industry groups: The American Plastics Council, and The Society of the Plastics Industry.

- 1. Plastics that go into curbside recycling bins get recycled.**
FACT: Most do not.
- 2. Curbside collection reduces the amount of plastic land filled.**
FACT: It does not.
- 3. Packaging resins are made from petroleum refinery waste.**
FACT: Nearly all are made from virgin petroleum and natural gas.
- 4. Plastic recyclers promote its recyclability.**
FACT: Plastic resin pellet producers pay for recycling ads to promote the sale of plastics.
- 5. Using plastic containers conserves energy.**
FACT: Most of the energy costs of plastic are incurred by the manufacturer. Virgin glass uses an equal amount of energy, while recycled glass uses far less than either virgin plastic or virgin glass.
- 6. Our choice is limited to recycling or wasting.**
FACT: Source reduction is key, and quite simple.



The History of Plastics

| 1869 | 1909 | 1939 | 1946 | 1955 | 1957 | 1983 | 2000 |
|---|---|--|---|---|---|---|--|
| John Wesley Hyatt invents celluloid, the first plastic product given a trade name | Bakelite is introduced to the Chemist Club in NY as the “first thermoset plastic,” meaning once set, it was set for life. | Nylon stockings debut at the World’s Fair. | Earl S. Tupper produces a 7-ounce polyethylene tumbler, the first of many products available from Tupperware Home Parties, Inc. | The Corvette is the first car to use plastic for body panels. | Monsanto’s House of Tomorrow, completely made of plastic, opens at Disneyland. The Hula Hoop creates a surge in demand for polyethylene | Microwave ovens open up a new market for plastic packaging. | In the U.S., pre-production plastic reaches 100 billion pounds of virgin resin pellets per year. |

What we do know about the North Pacific Ocean.

Scientists have been studying the problem of plastic floating pollution in the area since the 1970’s. These are some of their findings over the last few decades:

- In a 1999 study, the North Pacific Central Gyre was found to contain six pounds of plastic for each pound of surface zooplankton.
- The results of studies done in the 1980’s indicate that the quantity of plastic has tripled in the last ten years from maximum densities of 320,000 particles per square kilometer to 1 million particles per square kilometer.
- The filter-feeding animals in this area, mucous web feeding jellies and salps, were found to be heavily impacted by plastic fragments. The smaller the fragments, the fewer of them were found to be free floating, indicating that filter feeders had caught them.
- Filter feeders are at the lower end of the food chain, and fifty species of fish and many turtles are known to eat them, thus accumulating plastic in their stomachs.
- Plastic materials accumulate and concentrate organic chemicals and environmental pollutants up to one million times their concentration in the surrounding sea water. Many of these chemicals are called “endocrine disruptors”, and can be released when the plastics are ingested. The endocrine system produces hormones in humans and animals.
 - Hormones are amazingly potent. Estradiol, the body’s key estrogen hormone, operates at a concentration in the part per trillion range. One part per trillion is equivalent to one drop of water in 660 rail tank cars – a train 6 miles long.
 - Effects of hormone disruption on humans run the gamut from enlarged prostates and cancer to early puberty in young girls, even mental retardation and propensity to violence. In fish it can cause males to become female or fail to produce sperm.

For more information regarding this issue please view our video “Synthetic Sea: Plastics in the Open Ocean” available from AMRF for \$20 per copy.

Authors

Miwa Tamanaha, Santa Monica Bay Restoration Foundation.
Captain Charles Moore, Algalita Marine Research Foundation

Contributors

Anthony Andrady Ph.D., Research Triangle Institute
Michael Baker Ph.D., UCSD Department of Medicine
Kathy Cousins, Western Pacific Regional Fishery Management Council
Curtis Ebbesmeyer Ph.D., Beachcombers’ and Oceanographers’ International Association
Paul Goettlich, Berkeley Ecology Center Plastic Task Force
Robb Hamilton, Ornithologist
W. James Ingram, Jr., Alaska Fisheries Science Center, NOAA
Marilyn Judson, Pelagic Plastic Analysis Volunteer Coordinator
Tim Krupnik, Berkeley Ecology Center Plastic Task Force
Bill Macdonald, Bill Macdonald Productions
Shelly Moore, Southern California Coastal Water Research Project
Mark Spalding, International Environmental Policy and Law
Hideshige Takada Ph.D., Tokyo University of Agriculture and Technology
Tracy Williams, Aquarium of the Pacific, Long Beach, CA
Ann Zellers, Algalita Marine Research Foundation

Literature Cited

Day et al., 1985; Ryan, 1988a, 1990 (1992) Plastic Pellets in the Aquatic Environment Sources and Recommendations *EPA Final Report*, p. 31
Thorne-Miller, B. (November 1998) Island Press. *The Living Ocean: Understanding and Protecting Marine Biodiversity*, 2nd Edition
Twiss, J.R. and Reeves, R.R. ed. (September 1999) Conservation and Management of Marine Mammals. *Smithsonian Institution Press*, pp. 342-47
Various. (1993) Characterization of Municipal Solid Waste in the United States 1994 Update. *Environmental Protection Agency*.
Wallace, N. (1985) Debris entanglement in the marine environment: A review, pp. 259-277

